

# Neutrino and it's lepton

G.Quznetsov\*  
Chelyabinsk State University

November 10, 2008

## Abstract

In this paper I cite p.p. 100-117 of book G. Quznetsov, *Probabilistic Treatment of Gauge Theories*, in series Contemporary Fundamental Physics, ed. V. Dvoeglazov, Nova Sci. Publ., NY (2007). There I research a bound between neutrino and it's lepton.

Let  $\mathfrak{S}_{e\nu}$  be the unitary space, spanned by the following basis:

$$\mathbf{J}_{e\nu} := \left\langle \begin{array}{l} \frac{\hbar}{2\pi c} \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} (\cosh(\frac{\hbar}{c}n_0x_4) + \sinh(\frac{\hbar}{c}n_0x_4)) \epsilon_1, \\ \frac{\hbar}{2\pi c} \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} (\cosh(\frac{\hbar}{c}n_0x_4) + \sinh(\frac{\hbar}{c}n_0x_4)) \epsilon_2, \\ \frac{\hbar}{2\pi c} \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} (\cosh(\frac{\hbar}{c}n_0x_4) - \sinh(\frac{\hbar}{c}n_0x_4)) \epsilon_3, \\ \frac{\hbar}{2\pi c} \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} (\cosh(\frac{\hbar}{c}n_0x_4) - \sinh(\frac{\hbar}{c}n_0x_4)) \epsilon_4, \\ \frac{\hbar}{2\pi c} \exp(-i\frac{\hbar}{c}(n_0x_5)) \epsilon_1, \frac{\hbar}{2\pi c} \exp(-i\frac{\hbar}{c}(n_0x_5)) \epsilon_2, \\ \frac{\hbar}{2\pi c} \exp(-i\frac{\hbar}{c}(n_0x_5)) \epsilon_3, \frac{\hbar}{2\pi c} \exp(-i\frac{\hbar}{c}(n_0x_5)) \epsilon_4 \end{array} \right\rangle. \quad (1)$$

Let  $\mathfrak{S}_e$  be a subspace of the space  $\mathfrak{S}_{e\nu}$  such that if  $\tilde{\varphi} \in \mathfrak{S}_e$  then  $\tilde{\varphi}$  has got the following shape:

$$\tilde{\varphi}(t, \mathbf{x}, x_5, x_4) = \exp\left(-i\frac{\hbar}{c}n_0x_5\right) \sum_{k=1}^4 f_k(t, \mathbf{x}, n_0, 0) \epsilon_k$$

That is  $\tilde{\varphi}$  has got the following matrix in the basis  $\mathbf{J}_{e\nu}$ :

$$\tilde{\varphi} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ f_1 \\ f_2 \\ f_3 \\ f_4 \end{bmatrix}. \quad (2)$$

---

\*e-mails: gunn@mail.ru, quznets@yahoo.com

Let us consider the following Hamiltonian on  $\mathfrak{S}_e$ :

$$\begin{aligned}\widehat{H}_{0,4} &:= c \left( \sum_{r=1}^3 \beta^{[r]} i \partial_r + \gamma^{[0]} i \partial_5 + \beta^{[4]} i \partial_4 \right) : \\ \widehat{H}_{0,4} \tilde{\varphi} &= c \left( \sum_{r=1}^3 \beta^{[r]} i \partial_r + \gamma^{[0]} i \partial_5 + \beta^{[4]} i \partial_4 \right) \tilde{\varphi} = \\ &= \sum_{r=1}^3 \beta^{[r]} c i \partial_r \tilde{\varphi} + \\ &+ \gamma^{[0]} c i \partial_5 \exp \left( -i \frac{\hbar}{c} n_0 x_5 \right) \sum_{k=1}^4 f_k(t, \mathbf{x}, n_0, 0) \epsilon_k + \\ &+ \beta^{[4]} c i \partial_4 \exp \left( -i \frac{\hbar}{c} n_0 x_5 \right) \sum_{k=1}^4 f_k(t, \mathbf{x}, n_0, 0) \epsilon_k = \\ &= \sum_{r=1}^3 \beta^{[r]} c i \partial_r \tilde{\varphi} + \\ &+ \gamma^{[0]} c i \left( -i \frac{\hbar}{c} n_0 \right) \exp \left( -i \frac{\hbar}{c} n_0 x_5 \right) \sum_{k=1}^4 f_k(t, \mathbf{x}, n_0, 0) \epsilon_k + \\ &+ 0 = \\ &= \sum_{r=1}^3 \beta^{[r]} c i \partial_r \tilde{\varphi} + \hbar n_0 \gamma^{[0]} \exp \left( -i \frac{\hbar}{c} n_0 x_5 \right) \sum_{k=1}^4 f_k(t, \mathbf{x}, n_0, 0) \epsilon_k = \\ &= \sum_{r=1}^3 \beta^{[r]} c i \partial_r \tilde{\varphi} + \hbar n_0 \gamma^{[0]} \tilde{\varphi}.\end{aligned}\tag{3}$$

Hence on this space:

$$\widehat{H}_{0,4} = \widehat{H}_0 := c \sum_{r=1}^3 \beta^{[r]} i \partial_r + \hbar n_0 \gamma^{[0]}.\tag{4}$$

Let  $\mathfrak{S}_o$  be a subspace of the space  $\mathfrak{S}_{e\nu}$  such that if  $\tilde{\varphi}_o \in \mathfrak{S}_o$  then  $\tilde{\varphi}_o = \ell_o \tilde{\varphi} [1]$  and  $\tilde{\varphi} \in \mathfrak{S}_e$ , and if  $\tilde{\varphi} \in \mathfrak{S}_e$  then  $(\ell_o \tilde{\varphi}) \in \mathfrak{S}_o$ . If  $\tilde{\varphi}_o = \ell_o \tilde{\varphi}$  then in the basis  $\mathbf{J}_{e\nu}$ :

$$\tilde{\varphi}_o = \frac{1}{2\sqrt{(1-a^2)}} \begin{bmatrix} -(-q+ic) f_1 \\ -(-q+ic) f_2 \\ -(-q+ic) f_3 \\ -(-q+ic) f_4 \\ -\left(-\sqrt{(1-a^2)}+b\right) f_1 \\ -\left(-\sqrt{(1-a^2)}+b\right) f_2 \\ -\left(-\sqrt{(1-a^2)}+b\right) f_3 \\ -\left(-\sqrt{(1-a^2)}+b\right) f_4 \end{bmatrix}.$$

Let us consider a Hamiltonian  $\widehat{H}_{0,4}$  mode of behavior on the space  $\mathfrak{S}_o$ :

Hence

$$\begin{aligned}\widehat{H}_{0,4} \tilde{\varphi}_o &= c \sum_{r=1}^3 \beta^{[r]} i \partial_r \tilde{\varphi}_o + \\ &+ \gamma^{[0]} i c \frac{(q-ic)}{2\sqrt{(1-a^2)}} \frac{\hbar}{2\pi c} \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} \cdot \\ &\cdot \left( \partial_5 \left( \begin{aligned} &f_1 \left( \cosh \left( \frac{\hbar}{c} n_0 x_4 \right) + \sinh \left( \frac{\hbar}{c} n_0 x_4 \right) \right) \epsilon_1 + \\ &+ f_2 \left( \cosh \left( \frac{\hbar}{c} n_0 x_4 \right) + \sinh \left( \frac{\hbar}{c} n_0 x_4 \right) \right) \epsilon_2 + \\ &+ f_3 \left( \cosh \left( \frac{\hbar}{c} n_0 x_4 \right) - \sinh \left( \frac{\hbar}{c} n_0 x_4 \right) \right) \epsilon_3 + \\ &+ f_4 \left( \cosh \left( \frac{\hbar}{c} n_0 x_4 \right) - \sinh \left( \frac{\hbar}{c} n_0 x_4 \right) \right) \epsilon_4 + \end{aligned} \right) + \right. \\ &\left. + \left( \sqrt{(1-a^2)} - b \right) \frac{\hbar}{2\pi c} \partial_5 \exp \left( -i \frac{\hbar}{c} (n_0 x_5) \right) \cdot \right. \\ &\quad \cdot (f_1 \epsilon_1 + f_2 \epsilon_2 + f_3 \epsilon_3 + f_4 \epsilon_4) \end{aligned}$$

$$\begin{aligned}
& +\beta^{[4]}\mathrm{i}c\frac{(q-ic)}{2\sqrt{(1-a^2)}}\frac{\hbar}{2\pi c}\sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}}. \\
& \cdot \left( \partial_4 \left( \begin{aligned} & f_1 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_1 + \\ & + f_2 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_2 + \\ & + f_3 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_3 + \\ & + f_4 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_4 + \end{aligned} \right) + \right. \\
& \left. + \left( \sqrt{(1-a^2)} - b \right) \partial_4 \frac{\hbar}{2\pi c} \exp\left(-\mathrm{i}\frac{\hbar}{c}(n_0x_5)\right) \cdot \right. \\
& \quad \cdot (f_1\epsilon_1 + f_2\epsilon_2 + f_3\epsilon_3 + f_4\epsilon_4) \left. \right)
\end{aligned}$$

$$\begin{aligned}
& \widehat{H}_{0,4}\widetilde{\varphi}_0 = c \sum_{r=1}^3 \beta^{[r]}\mathrm{i}\partial_r\widetilde{\varphi}_0 + \\
& + \gamma^{[0]}\mathrm{i}c\frac{(q-ic)}{2\sqrt{(1-a^2)}}\frac{\hbar}{2\pi c}\sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}}. \\
& \cdot \left( \partial_5 \left( \begin{aligned} & f_1 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_1 + \\ & + f_2 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_2 + \\ & + f_3 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_3 + \\ & + f_4 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_4 + \end{aligned} \right) + \right. \\
& \left. + \left( \sqrt{(1-a^2)} - b \right) \frac{\hbar}{2\pi c} \partial_5 \exp\left(-\mathrm{i}\frac{\hbar}{c}(n_0x_5)\right) \cdot \right. \\
& \quad \cdot (f_1\epsilon_1 + f_2\epsilon_2 + f_3\epsilon_3 + f_4\epsilon_4) \left. \right) +
\end{aligned}$$

$$\begin{aligned}
& +\beta^{[4]}\mathrm{i}c\frac{(q-ic)}{2\sqrt{(1-a^2)}}\frac{\hbar}{2\pi c}\sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}}. \\
& \cdot \left( \partial_4 \left( \begin{aligned} & f_1 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_1 + \\ & + f_2 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_2 + \\ & + f_3 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_3 + \\ & + f_4 \left( \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_4 + \end{aligned} \right) + \right. \\
& \left. + \left( \sqrt{(1-a^2)} - b \right) \partial_4 \frac{\hbar}{2\pi c} \exp\left(-\mathrm{i}\frac{\hbar}{c}(n_0x_5)\right) \cdot \right. \\
& \quad \cdot (f_1\epsilon_1 + f_2\epsilon_2 + f_3\epsilon_3 + f_4\epsilon_4) \left. \right)
\end{aligned}$$

Therefore

$$\begin{aligned}
& \widehat{H}_{0,4}\widetilde{\varphi}_0 = c \sum_{r=1}^3 \beta^{[r]}\mathrm{i}\partial_r\widetilde{\varphi}_0 + \\
& + \gamma^{[0]}\mathrm{i}c\frac{\sqrt{1-a^2}-b}{2\sqrt{1-a^2}}. \\
& \cdot \left( 0 + \frac{\hbar}{2\pi c} \partial_5 \exp\left(-\mathrm{i}\frac{\hbar}{c}(n_0x_5)\right) (f_1\epsilon_1 + f_2\epsilon_2 + f_3\epsilon_3 + f_4\epsilon_4) \right) + \\
& + \beta^{[4]}\mathrm{i}c\frac{q-ic}{2\sqrt{1-a^2}}\frac{\hbar}{2\pi c}\sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}}. \\
& \cdot \left( \left( \begin{aligned} & f_1 \left( \partial_4 \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \partial_4 \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_1 + \\ & + f_2 \left( \partial_4 \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \partial_4 \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_2 + \\ & + f_3 \left( \partial_4 \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \partial_4 \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_3 + \\ & + f_4 \left( \partial_4 \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \partial_4 \sinh\left(\frac{\hbar}{c}n_0x_4\right) \right) \epsilon_4 + \end{aligned} \right) + \right. \\
& \quad + 0 \left. \right)
\end{aligned}$$

Hence

$$\begin{aligned}
\hat{H}_{0,4}\tilde{\varphi}_o &= c \sum_{r=1}^3 \beta^{[r]} i \partial_r \tilde{\varphi}_o + \\
&+ \gamma^{[0]} i c \left( -i \frac{h}{c} n_0 \right) \frac{\sqrt{1-a^2}-b}{2\sqrt{1-a^2}} \frac{h}{2\pi c} \exp \left( -i \frac{h}{c} n_0 x_5 \right) \cdot \\
&\cdot (f_1 \epsilon_1 + f_2 \epsilon_2 + f_3 \epsilon_3 + f_4 \epsilon_4) + \\
&+ \beta^{[4]} i c \frac{h}{c} n_0 \frac{q-ic}{2\sqrt{1-a^2}} \frac{h}{2\pi c} \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} \cdot \\
&\cdot \begin{pmatrix} f_1 \left( \sinh \left( \frac{h}{c} n_0 x_4 \right) + \cosh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_1 + \\ + f_2 \left( \sinh \left( \frac{h}{c} n_0 x_4 \right) + \cosh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_2 + \\ + f_3 \left( \sinh \left( \frac{h}{c} n_0 x_4 \right) - \cosh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_3 + \\ + f_4 \left( \sinh \left( \frac{h}{c} n_0 x_4 \right) - \cosh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_4 + \end{pmatrix}.
\end{aligned}$$

Therefore

$$\begin{aligned}
\hat{H}_{0,4}\tilde{\varphi}_o &= c \sum_{r=1}^3 \beta^{[r]} i \partial_r \tilde{\varphi}_o + \\
&+ h n_0 \gamma^{[0]} \frac{\sqrt{1-a^2}-b}{2\sqrt{1-a^2}} \frac{h}{2\pi c} \exp \left( -i \frac{h}{c} n_0 x_5 \right) \cdot \\
&\cdot (f_1 \epsilon_1 + f_2 \epsilon_2 + f_3 \epsilon_3 + f_4 \epsilon_4) + \\
&+ h n_0 \beta^{[4]} i \frac{q-ic}{2\sqrt{1-a^2}} \frac{h}{2\pi c} \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} \cdot \\
&\cdot \begin{pmatrix} f_1 \left( \cosh \left( \frac{h}{c} n_0 x_4 \right) + \sinh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_1 + \\ + f_2 \left( \cosh \left( \frac{h}{c} n_0 x_4 \right) + \sinh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_2 - \\ - f_3 \left( \cosh \left( \frac{h}{c} n_0 x_4 \right) - \sinh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_3 - \\ - f_4 \left( \cosh \left( \frac{h}{c} n_0 x_4 \right) - \sinh \left( \frac{h}{c} n_0 x_4 \right) \right) \epsilon_4 + \end{pmatrix}.
\end{aligned}$$

Hence in the basis  $\mathbf{J}_{e\nu}$ :

$$\begin{aligned}
\hat{H}_{0,4}\tilde{\varphi}_o &= c \sum_{r=1}^3 \beta^{[r]} i \partial_r \tilde{\varphi}_o + h n_0 \cdot \\
&\cdot \left( \gamma^{[0]} \frac{\sqrt{1-a^2}-b}{2\sqrt{1-a^2}} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ f_1 \\ f_2 \\ f_3 \\ f_4 \end{bmatrix} + \beta^{[4]} i \frac{q-ic}{2\sqrt{1-a^2}} \begin{bmatrix} f_1 \\ f_2 \\ -f_3 \\ -f_4 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right) = \\
&= c \sum_{r=1}^3 \beta^{[r]} i \partial_r \tilde{\varphi}_o + h n_0 \cdot \\
&\cdot \left( \gamma^{[0]} \frac{\sqrt{1-a^2}-b}{2\sqrt{1-a^2}} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ f_1 \\ f_2 \\ f_3 \\ f_4 \end{bmatrix} + \beta^{[4]} i \frac{q-ic}{2\sqrt{1-a^2}} \gamma^{[5]} \begin{bmatrix} f_1 \\ f_2 \\ f_3 \\ f_4 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right).
\end{aligned}$$

with

$$\gamma^{[5]} := \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}.$$

Since

$$\beta^{[4]}\mathbf{i}\gamma^{[5]} = \gamma^{[0]}$$

then

$$\begin{aligned} \widehat{H}_{0,4}\widetilde{\varphi}_o &= \mathbf{c} \sum_{r=1}^3 \beta^{[r]}\mathbf{i}\partial_r\widetilde{\varphi}_o + \mathbf{h}n_0 \cdot \\ &\cdot \left( \gamma^{[0]} \frac{1}{2\sqrt{1-a^2}} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ (\sqrt{1-a^2}-b)f_1 \\ (\sqrt{1-a^2}-b)f_2 \\ (\sqrt{1-a^2}-b)f_3 \\ (\sqrt{1-a^2}-b)f_4 \end{bmatrix} + \gamma^{[0]} \frac{1}{2\sqrt{1-a^2}} \begin{bmatrix} (q-ic)f_1 \\ (q-ic)f_2 \\ (q-ic)f_3 \\ (q-ic)f_4 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right). \end{aligned}$$

Therefore

$$\widehat{H}_{0,4}\widetilde{\varphi}_o = \mathbf{c} \sum_{r=1}^3 \beta^{[r]}\mathbf{i}\partial_r\widetilde{\varphi}_o + \mathbf{h}n_0\gamma^{[0]} \frac{1}{2\sqrt{1-a^2}} \begin{pmatrix} -(-q+ic)f_1, \\ -(-q+ic)f_2, \\ -(-q+ic)f_3, \\ -(-q+ic)f_4, \\ -(-\sqrt{1-a^2}+b)f_1, \\ -(-\sqrt{1-a^2}+b)f_2, \\ -(-\sqrt{1-a^2}+b)f_3, \\ -(-\sqrt{1-a^2}+b)f_4 \end{pmatrix}.$$

Hence

$$\widehat{H}_{0,4}\widetilde{\varphi}_o = \mathbf{c} \sum_{r=1}^3 \beta^{[r]}\mathbf{i}\partial_r\widetilde{\varphi}_o + \mathbf{h}n_0\gamma^{[0]}\widetilde{\varphi}_o.$$

Thus in the space  $\mathfrak{S}_e$ :

$$\widehat{H}_{0,4} = \widehat{H}_0 = \mathbf{c} \sum_{r=1}^3 \beta^{[r]}\mathbf{i}\partial_r + \mathbf{h}n_0\gamma^{[0]},$$

too.

Let  $\mathfrak{S}_*$  be a subspace of the space  $\mathfrak{S}_{e\nu}$  such that if  $\widetilde{\varphi}_* \in \mathfrak{S}_*$  then  $\widetilde{\varphi}_* = \ell_* \widetilde{\varphi} [1]$  and  $\widetilde{\varphi} \in \mathfrak{S}_e$ , and if  $\widetilde{\varphi} \in \mathfrak{S}_e$  then  $(\ell_* \widetilde{\varphi}) \in \mathfrak{S}_*$ . If  $\widetilde{\varphi}_* = \ell_* \widetilde{\varphi} (2)$  then in the basis  $\mathbf{J}_{e\nu}$ :

$$\tilde{\varphi}_* = \frac{1}{2\sqrt{(1-a^2)}} \begin{bmatrix} (-q+ic) f_1 \\ (-q+ic) f_2 \\ (-q+ic) f_3 \\ (-q+ic) f_4 \\ (b+\sqrt{1-a^2}) f_1 \\ (b+\sqrt{1-a^2}) f_2 \\ (b+\sqrt{1-a^2}) f_3 \\ (b+\sqrt{1-a^2}) f_4 \end{bmatrix}.$$

Similarly to  $\tilde{\varphi}_0$  you can calculate that

$$\hat{H}_{0,4}\tilde{\varphi}_* = \hat{H}_0\tilde{\varphi}_* = c \sum_{r=1}^3 \beta^{[r]} i \partial_r \tilde{\varphi}_* + \hbar n_0 \gamma^{[0]} \tilde{\varphi}_*,$$

too.

Let

$$e_{1L}(\mathbf{k}) := \begin{bmatrix} \omega(\mathbf{k}) + n_0 + k_3 \\ k_1 + i k_2 \end{bmatrix}, \quad e_{1R}(\mathbf{k}) := \begin{bmatrix} \omega(\mathbf{k}) + n_0 - k_3 \\ -k_1 - i k_2 \end{bmatrix},$$

$$e_{2L}(\mathbf{k}) := \begin{bmatrix} k_1 - i k_2 \\ \omega(\mathbf{k}) + n_0 - k_3 \end{bmatrix}, \quad e_{2R}(\mathbf{k}) := \begin{bmatrix} -k_1 + i k_2 \\ \omega(\mathbf{k}) + n_0 + k_3 \end{bmatrix},$$

$$\begin{aligned} e_{3L}(\mathbf{k}) &:= -e_{1R}(\mathbf{k}), \quad e_{3R}(\mathbf{k}) := e_{1L}(\mathbf{k}), \\ e_{4L}(\mathbf{k}) &:= -e_{2R}(\mathbf{k}), \quad e_{4R}(\mathbf{k}) := e_{2L}(\mathbf{k}). \end{aligned}$$

with

$$\omega(\mathbf{k}) := \sqrt{n_0^2 + k_1^2 + k_2^2 + k_3^2}$$

( $n_0, k_1, k_2, k_3$  are real numbers).

In this case [2]:

$$e_s(\mathbf{k}) = \frac{1}{2\sqrt{\omega(\mathbf{k})(\omega(\mathbf{k}) + n_0)}} \begin{bmatrix} e_{sL}(\mathbf{k}) \\ e_{sR}(\mathbf{k}) \end{bmatrix}.$$

Let

$$\underline{e}_s(\mathbf{k}) := \begin{bmatrix} \vec{0}_4 \\ e_s(\mathbf{k}) \end{bmatrix}$$

here  $s \in \{1, 2, 3, 4\}$ .

And let [3]:

$$\begin{aligned} \underline{e}_{os}(\mathbf{k}) &:= \ell_o \underline{e}_s(\mathbf{k}) \\ &= \frac{1}{\sqrt{2(\sqrt{1-a^2}-b)}\sqrt{1-a^2}} \begin{bmatrix} (q-ic) e_s(\mathbf{k}) \\ (\sqrt{1-a^2}-b) e_s(\mathbf{k}) \end{bmatrix}, \end{aligned}$$

$$\begin{aligned} \underline{e}_{*s}(\mathbf{k}) &:= \ell_* \underline{e}_s(\mathbf{k}) \\ &= \frac{1}{\sqrt{2(\sqrt{1-a^2}+b)}\sqrt{1-a^2}} \begin{bmatrix} (-q+ic) e_s(\mathbf{k}) \\ (b+\sqrt{1-a^2}) e_s(\mathbf{k}) \end{bmatrix}. \end{aligned}$$

Denote

$$\hat{H}_0(\mathbf{k}) := \sum_{r=1}^3 \beta^{[r]} k_r = \begin{bmatrix} k_3 & k_1 - ik_2 & n_0 & 0 \\ k_1 + ik_2 & -k_3 & 0 & n_0 \\ n_0 & 0 & -k_3 & -k_1 + ik_2 \\ 0 & n_0 & -k_1 - ik_2 & k_3 \end{bmatrix}.$$

In that case

$$\begin{aligned} \hat{H}_0 \underline{e}_{o1}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right) &= \\ &= \hbar \hat{H}_0(\mathbf{k}) \underline{e}_{o1}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right) \\ &= \hbar \omega(\mathbf{k}) \underline{e}_{o1}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right). \end{aligned}$$

Therefore  $\underline{e}_{o1}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$  is an eigenvector of  $\hat{H}_0$  with the eigenvalue  $\hbar \omega(\mathbf{k})$ . Similarly you can calculate that

$\underline{e}_{o2}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$ ,  $\underline{e}_{*1}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$ ,  $\underline{e}_{*2}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$ , are eigenvectors of  $\hat{H}_0$  with the same eigenvalue, and

$\underline{e}_{o3}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$ ,  $\underline{e}_{o4}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$ ,

$\underline{e}_{*3}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$ ,  $\underline{e}_{*4}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$

are an eigenvectors of  $\hat{H}_0$  with the eigenvalue  $(-\hbar \omega(\mathbf{k}))$ .

Vectors  $\underline{e}_{os}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$ ,  $\underline{e}_{*s}(\mathbf{k}) \left( \frac{\hbar}{2\pi c} \right)^{\frac{3}{2}} \exp\left( i \frac{\hbar}{c} \right)$  with  $s \in \{1, 2, 3, 4\}$  form an orthonormalized basis in the space  $\mathfrak{S}_{e\nu}(1)$  and

$$\sum_{s=1}^4 (\underline{e}_{*s,r}^*(\mathbf{k}) \underline{e}_{*s,r'}(\mathbf{k}) + \underline{e}_{os,r}^*(\mathbf{k}) \underline{e}_{os,r'}(\mathbf{k})) = \delta_{r,r'} \quad (5)$$

for  $r, r' \in \{1, 2, 3, 4, 5, 6, 7, 8\}$ .

Let [3]

$$\begin{aligned} \underline{e}'_{*s}(\mathbf{k}) &:= U^{(-)} \underline{e}_{*s}(\mathbf{k}) \\ &= \frac{1}{\sqrt{2(\sqrt{1-a^2}+b)}\sqrt{1-a^2}} \frac{1}{2\sqrt{\omega(\mathbf{k})(\omega(\mathbf{k})+n_0)}} \\ &\quad \cdot \begin{bmatrix} (a-i\sqrt{1-a^2})(-q+ic)e_{sL}(\mathbf{k}) \\ (-q+ic)e_{sR}(\mathbf{k}) \\ (a-i\sqrt{1-a^2})(\sqrt{1-a^2}+b)e_{sL}(\mathbf{k}) \\ (\sqrt{1-a^2}+b)e_{sR}(\mathbf{k}) \end{bmatrix} \end{aligned}$$

and

$$\begin{aligned} \underline{e}'_{os}(\mathbf{k}) &:= U^{(-)} \underline{e}_{os}(\mathbf{k}) \\ &= \frac{1}{\sqrt{2(\sqrt{1-a^2}-b)}\sqrt{1-a^2}} \frac{1}{2\sqrt{\omega(\mathbf{k})(\omega(\mathbf{k})+n_0)}} \\ &\quad \cdot \begin{bmatrix} (a+i\sqrt{1-a^2})(q-ic)e_{sL}(\mathbf{k}) \\ (q-ic)e_{sR}(\mathbf{k}) \\ (a+i\sqrt{1-a^2})(\sqrt{1-a^2}-b)e_{sL}(\mathbf{k}) \\ (\sqrt{1-a^2}-b)e_{sR}(\mathbf{k}) \end{bmatrix}. \end{aligned}$$

For these vectors:

$$\sum_{r=1}^4 (\underline{e}'_{*r,j}(\mathbf{k}) \underline{e}'_{*r,j'}(\mathbf{k}) + \underline{e}'_{or,j}(\mathbf{k}) \underline{e}'_{or,j'}(\mathbf{k})) = \delta_{j,j'}$$

and since  $U^{(-)\dagger}U^{(-)} = 1_8$  then  $\underline{e}'_{os}(\mathbf{k})$  and  $\underline{e}'_{*s}(\mathbf{k})$  form an orthonormalized basis in the space  $\mathfrak{S}_{e\nu}$ , too.

Let

$$\underline{e}'_r(\mathbf{k}) := U^{(-)} \underline{e}_r(\mathbf{k}) = \frac{1}{2\sqrt{\omega(\mathbf{k})(\omega(\mathbf{k})+n_0)}} \begin{bmatrix} (c+iq)e_{rL}(\mathbf{k}) \\ \vec{0}_2 \\ (a-ib)e_{rL}(\mathbf{k}) \\ e_{rR}(\mathbf{k}) \end{bmatrix}. \quad (6)$$

In that case:

$$\underline{e}'_r(\mathbf{k}) = \frac{1}{\sqrt{2}} \left( \sqrt{1 - \frac{b}{\sqrt{1-a^2}}} \underline{e}'_{or}(\mathbf{k}) + \sqrt{1 + \frac{b}{\sqrt{1-a^2}}} \underline{e}'_{*r}(\mathbf{k}) \right).$$

Let for  $j, j' \in \{1, 2, 3, 4, 5, 6, 7, 8\}$  [4]:



$$\begin{aligned}\left\{\psi_{j'}^\dagger(\mathbf{y}), \psi_j(\mathbf{x})\right\} &= \delta(\mathbf{y} - \mathbf{x}) \delta_{j',j} \widehat{1}, \\ \left\{\psi_{j'}^\dagger(\mathbf{y}), \psi_j^\dagger(\mathbf{x})\right\} &= \widehat{0} = \{\psi_{j'}(\mathbf{y}), \psi_j(\mathbf{x})\}\end{aligned}$$

and let

$$\begin{aligned}b_{or,\mathbf{k}} &:= \left(\frac{\hbar}{2\pi c}\right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{j'=1}^8 \mathcal{E}_{or,j'}^*(\mathbf{k}) \psi_{j'}(\mathbf{x}), \\ b_{*r,\mathbf{k}} &:= \left(\frac{\hbar}{2\pi c}\right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{j'=1}^8 \mathcal{E}_{*r,j'}^*(\mathbf{k}) \psi_{j'}(\mathbf{x}).\end{aligned}$$

In that case:

$$\begin{aligned}& \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( \sum_{r=1}^4 \mathcal{E}_{or,j}(\mathbf{k}) b_{or,\mathbf{k}} + \sum_{r=1}^4 \mathcal{E}_{*r,j}(\mathbf{k}) b_{*r,\mathbf{k}} \right) \\ &= \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( \begin{aligned} & \sum_{r=1}^4 \mathcal{E}_{or,j}(\mathbf{k}) \left(\frac{\hbar}{2\pi c}\right)^3 \cdot \\ & \cdot \int_{(\Omega)} d\mathbf{x}' \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}'} \sum_{j'=1}^8 \mathcal{E}_{or,j'}^*(\mathbf{k}) \psi_{j'}(\mathbf{x}') \\ & + \sum_{r=1}^4 \mathcal{E}_{*r,j}(\mathbf{k}) \left(\frac{\hbar}{2\pi c}\right)^3 \cdot \\ & \cdot \int_{(\Omega)} d\mathbf{x}' \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}'} \sum_{j'=1}^8 \mathcal{E}_{*r,j'}^*(\mathbf{k}) \psi_{j'}(\mathbf{x}') \end{aligned} \right) \\ &= \left(\frac{\hbar}{2\pi c}\right)^3 \sum_{\mathbf{k}} \int_{(\Omega)} d\mathbf{x}' \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}'} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \cdot \\ & \cdot \sum_{r=1}^4 \sum_{j'=1}^8 (\mathcal{E}_{or,j}(\mathbf{k}) \mathcal{E}_{or,j'}^*(\mathbf{k}) + \mathcal{E}_{*r,j}(\mathbf{k}) \mathcal{E}_{*r,j'}^*(\mathbf{k})) \psi_{j'}(\mathbf{x}').\end{aligned}$$

In accordance with (5):

$$\begin{aligned}& \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( \sum_{r=1}^4 \mathcal{E}_{or,j}(\mathbf{k}) b_{or,\mathbf{k}} + \sum_{r=1}^4 \mathcal{E}_{*r,j}(\mathbf{k}) b_{*r,\mathbf{k}} \right) \\ &= \left(\frac{\hbar}{2\pi c}\right)^3 \int_{(\Omega)} d\mathbf{x}' \cdot \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}(\mathbf{x}-\mathbf{x}')} \sum_{j'=1}^8 \delta_{j,j'} \psi_{j'}(\mathbf{x}').\end{aligned}$$

Hence since

$$\sum_{\mathbf{k}} e^{i\frac{\hbar}{c}\mathbf{k}(\mathbf{x}'-\mathbf{x})} = \left(\frac{h}{2\pi c}\right)^3 \delta(\mathbf{x}' - \mathbf{x})$$

and according properties of  $\delta$ :

$$\begin{aligned}
& \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( \sum_{r=1}^4 \underline{e}_{or,j}(\mathbf{k}) b_{or,\mathbf{k}} + \sum_{r=1}^4 \underline{e}_{*r,j}(\mathbf{k}) b_{*r,\mathbf{k}} \right) \\
&= \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x}' \cdot \left( \frac{\hbar}{2\pi c} \right)^3 \delta(\mathbf{x}' - \mathbf{x}) \psi_j(\mathbf{x}') \\
&= \int_{(\Omega)} d\mathbf{x}' \cdot \delta(\mathbf{x}' - \mathbf{x}) \psi_j(\mathbf{x}') = \psi_j(\mathbf{x}).
\end{aligned}$$

Thus:

$$\boxed{\sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( \sum_{r=1}^4 \underline{e}_{or,j}(\mathbf{k}) b_{or,\mathbf{k}} + \sum_{r=1}^4 \underline{e}_{*r,j}(\mathbf{k}) b_{*r,\mathbf{k}} \right) = \psi_j(\mathbf{x}).} \quad (7)$$

Let

$$\begin{aligned}
& \psi(\mathbf{x}) := \frac{\hbar}{2\pi c} \cdot \\
& \cdot \left( \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} \left( \begin{pmatrix} \cosh\left(\frac{\hbar}{c}n_0x_4\right) + \\ + \sinh\left(\frac{\hbar}{c}n_0x_4\right) \end{pmatrix} \sum_{r=1}^2 \psi_r(\mathbf{x}) \epsilon_r + \right. \right. \\
& \quad \left. \left. + \begin{pmatrix} \cosh\left(\frac{\hbar}{c}n_0x_4\right) - \\ - \sinh\left(\frac{\hbar}{c}n_0x_4\right) \end{pmatrix} \sum_{r=3}^4 \psi_r(\mathbf{x}) \epsilon_r \right) + \right. \\
& \quad \left. + \exp\left(-i\frac{\hbar}{c}(n_0x_4)\right) \sum_{r=1}^4 \psi_{r+4}(\mathbf{x}) \epsilon_r \right). \quad (8)
\end{aligned}$$

That is in the basis  $\mathbf{J}_{e\nu}(1)$ :

$$\psi(\mathbf{x}) = \begin{bmatrix} \psi_1(\mathbf{x}) \\ \psi_2(\mathbf{x}) \\ \psi_3(\mathbf{x}) \\ \psi_4(\mathbf{x}) \\ \psi_5(\mathbf{x}) \\ \psi_6(\mathbf{x}) \\ \psi_7(\mathbf{x}) \\ \psi_8(\mathbf{x}) \end{bmatrix}.$$

That is in this basis:

$$\begin{aligned}
b_{or,\mathbf{k}} &:= \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \underline{e}_{or,j}^\dagger(\mathbf{k}) \psi(\mathbf{x}), \\
b_{*r,\mathbf{k}} &:= \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \underline{e}_{*r,j}^\dagger(\mathbf{k}) \psi(\mathbf{x}).
\end{aligned}$$

Let

$$\psi'(\mathbf{x}) := U^{(-)} \psi(\mathbf{x}).$$

In that case:

$$b'_{or,\mathbf{k}} := \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \underline{e}'_{or,j}{}^\dagger(\mathbf{k}) \psi'(\mathbf{x}),$$

$$b'_{*r,\mathbf{k}} := \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \underline{e}'_{*r,j}{}^\dagger(\mathbf{k}) \psi'(\mathbf{x}).$$

Hence:

$$b'_{or,\mathbf{k}} = \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( U^{(-)} \underline{e}_{or,j}(\mathbf{k}) \right)^\dagger \left( U^{(-)} \psi(\mathbf{x}) \right),$$

$$b'_{*r,\mathbf{k}} = \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( U^{(-)} \underline{e}_{*r,j}(\mathbf{k}) \right)^\dagger \left( U^{(-)} \psi(\mathbf{x}) \right).$$

Since  $U^{(-)\dagger}U^{(-)} = 1_8$  then

$$b'_{or,\mathbf{k}} = \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \underline{e}_{or,j}{}^\dagger(\mathbf{k}) \psi(\mathbf{x}),$$

$$b'_{*r,\mathbf{k}} = \left( \frac{\hbar}{2\pi c} \right)^3 \int_{(\Omega)} d\mathbf{x} \cdot e^{i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \underline{e}_{*r,j}{}^\dagger(\mathbf{k}) \psi(\mathbf{x}).$$

That is:

$$b'_{or,\mathbf{k}} = b_{or,\mathbf{k}} \text{ and } b'_{*r,\mathbf{k}} = b_{*r,\mathbf{k}}.$$

And from (7):

$$\psi'_j(\mathbf{x}) = \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{r=1}^4 \left( \underline{e}'_{or,j}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r,j}(\mathbf{k}) b_{*r,\mathbf{k}} \right). \quad (9)$$

For the operators  $b_{or,\mathbf{k}}$  and  $b_{*r,\mathbf{k}}$ :

$$\begin{aligned} \left\{ b_{or',\mathbf{k}'}^\dagger, b_{or,\mathbf{k}} \right\} &= \left( \frac{\hbar}{2\pi c} \right)^3 \delta_{r,r'} \delta_{\mathbf{k},\mathbf{k}'} \widehat{1}, \\ \left\{ b_{*r',\mathbf{k}'}^\dagger, b_{*r,\mathbf{k}} \right\} &= \left( \frac{\hbar}{2\pi c} \right)^3 \delta_{r,r'} \delta_{\mathbf{k},\mathbf{k}'} \widehat{1}, \\ \left\{ b_{or',\mathbf{k}'}^\dagger, b_{*r,\mathbf{k}} \right\} &= \widehat{0}, \\ \left\{ b_{*r',\mathbf{k}'}^\dagger, b_{or,\mathbf{k}} \right\} &= \widehat{0}, \\ \left\{ b_{or',\mathbf{k}'}^\dagger, b_{or,\mathbf{k}} \right\} &= \widehat{0}, \\ \left\{ b_{*r',\mathbf{k}'}^\dagger, b_{*r,\mathbf{k}} \right\} &= \widehat{0}, \\ \left\{ b_{or',\mathbf{k}'}^\dagger, b_{or,\mathbf{k}} \right\} &= \widehat{0}, \\ \left\{ b_{*r',\mathbf{k}'}^\dagger, b_{*r,\mathbf{k}} \right\} &= \widehat{0}, \\ \left\{ b_{*r',\mathbf{k}'}^\dagger, b_{or,\mathbf{k}} \right\} &= \widehat{0}. \end{aligned}$$

(10)

Let

$$b_{r,\mathbf{k}} := \sqrt{2} (1 - a^2)^{\frac{1}{4}} \left( \frac{1}{\sqrt{\sqrt{1 - a^2} - b}} b_{or,\mathbf{k}} + \frac{1}{\sqrt{\sqrt{1 - a^2} + b}} b_{*r,\mathbf{k}} \right).$$

In that case:

$$\begin{aligned} & \underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}} = \\ &= \frac{1}{\sqrt{2}} \left( \sqrt{1 - \frac{b}{\sqrt{1 - a^2}}} \underline{e}'_{or}(\mathbf{k}) + \sqrt{1 + \frac{b}{\sqrt{1 - a^2}}} \underline{e}'_{*r}(\mathbf{k}) \right) b_{r,\mathbf{k}} \\ & \quad - \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \underline{e}'_{or}(\mathbf{k}) b_{*r,\mathbf{k}} - \sqrt{\frac{b + \sqrt{1 - a^2}}{b - \sqrt{1 - a^2}}} \underline{e}'_{*r}(\mathbf{k}) b_{or,\mathbf{k}} \end{aligned}$$

And from (6):

$$\begin{aligned} & \underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}} \\ &= \underline{e}'_r(\mathbf{k}) b_{r,\mathbf{k}} \\ & \quad - \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \underline{e}'_{or}(\mathbf{k}) b_{*r,\mathbf{k}} \\ & \quad - \sqrt{\frac{b + \sqrt{1 - a^2}}{b - \sqrt{1 - a^2}}} \underline{e}'_{*r}(\mathbf{k}) b_{or,\mathbf{k}}. \end{aligned}$$

For  $b_{r,\mathbf{k}}$ :

$$\boxed{\begin{aligned} \{b_{r',\mathbf{k}'}, b_{r,\mathbf{k}}\} &= 4 \frac{b^2 + c^2 + q^2}{c^2 + q^2} \left( \frac{h}{2\pi c} \right)^3 \delta_{r,r'} \delta_{\mathbf{k},\mathbf{k}'} \widehat{1}, \\ \{b_{r',\mathbf{k}'}, b_{r,\mathbf{k}}^\dagger\} &= \widehat{0}, \\ \{b_{r',\mathbf{k}'}, b_{r,\mathbf{k}}\} &= \widehat{0}. \end{aligned}} \quad (11)$$

From (9):

$$\begin{aligned} \psi'_j(\mathbf{x}) &= \sum_{\mathbf{k}} e^{-i \frac{h}{c} \mathbf{k} \mathbf{x}} \sum_{r=1}^4 \underline{e}'_{r,j}(\mathbf{k}) b_{r,\mathbf{k}} \\ & \quad - \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} e^{-i \frac{h}{c} \mathbf{k} \mathbf{x}} \sum_{r=1}^4 \underline{e}'_{or,j}(\mathbf{k}) b_{*r,\mathbf{k}} \\ & \quad - \sqrt{\frac{b + \sqrt{1 - a^2}}{b - \sqrt{1 - a^2}}} \sum_{\mathbf{k}} e^{-i \frac{h}{c} \mathbf{k} \mathbf{x}} \sum_{r=1}^4 \underline{e}'_{*r,j}(\mathbf{k}) b_{or,\mathbf{k}}. \end{aligned}$$

Let:

$$\chi(\mathbf{x}) := \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{r=1}^4 \underline{e}'_r(\mathbf{k}) b_{r,\mathbf{k}}, \quad (12)$$

$$\chi_{*j}(\mathbf{x}) := \sqrt{\frac{b - \sqrt{1-a^2}}{b + \sqrt{1-a^2}}} \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{r=1}^4 \underline{e}'_{or,j}(\mathbf{k}) b_{*r,\mathbf{k}}$$

$$\chi_{oj}(\mathbf{x}) := \sqrt{\frac{b + \sqrt{1-a^2}}{b - \sqrt{1-a^2}}} \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{r=1}^4 \underline{e}'_{*r,j}(\mathbf{k}) b_{or,\mathbf{k}}.$$

In that case:

$$\psi'_j(\mathbf{x}) = \chi_j(\mathbf{x}) - \chi_{*j}(\mathbf{x}) - \chi_{oj}(\mathbf{x}).$$

Let

$$\hat{H}'_0 := U^{(-)} \hat{H}_0 U^{(-)\dagger}.$$

For this Hamiltonian:

$$\begin{aligned} & \int_{(\Omega)} d\mathbf{x} \cdot \chi_*^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \\ &= \int_{(\Omega)} d\mathbf{x} \cdot \sqrt{\frac{b - \sqrt{1-a^2}}{b + \sqrt{1-a^2}}} \sum_{\mathbf{k}'} e^{i\frac{\hbar}{c}\mathbf{k}'\mathbf{x}} \sum_{r'=1}^4 b_{*r',\mathbf{k}'}^\dagger \underline{e}_{or'}'^\dagger(\mathbf{k}') \cdot \\ & \quad \cdot \hat{H}'_0 \sum_{\mathbf{k}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{r=1}^4 (\underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}}) \\ &= \int_{(\Omega)} d\mathbf{x} \cdot \sqrt{\frac{b - \sqrt{1-a^2}}{b + \sqrt{1-a^2}}} \sum_{\mathbf{k}} \sum_{\mathbf{k}'} e^{i\frac{\hbar}{c}\mathbf{k}'\mathbf{x}} \sum_{r'=1}^4 b_{*r',\mathbf{k}'}^\dagger \underline{e}_{or'}'^\dagger(\mathbf{k}') \cdot \\ & \quad \cdot \hat{H}'_0 e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \left( \sum_{r=1}^2 (\underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}}) + \right. \\ & \quad \left. + \sum_{r=3}^4 (\underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}}) \right) \\ &= \int_{(\Omega)} d\mathbf{x} \cdot \sqrt{\frac{b - \sqrt{1-a^2}}{b + \sqrt{1-a^2}}} \sum_{\mathbf{k}} \sum_{\mathbf{k}'} e^{i\frac{\hbar}{c}\mathbf{k}'\mathbf{x}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{r'=1}^4 b_{*r',\mathbf{k}'}^\dagger \underline{e}_{or'}'^\dagger(\mathbf{k}') \cdot \\ & \quad \cdot \hbar \hat{H}'_0(\mathbf{k}) \left( \sum_{r=1}^2 (\underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}}) + \right. \\ & \quad \left. + \sum_{r=3}^4 (\underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}}) \right). \end{aligned}$$

Hence:

$$\begin{aligned}
& \int_{(\Omega)} d\mathbf{x} \cdot \chi_*^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \\
&= \int_{(\Omega)} d\mathbf{x} \cdot \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \sum_{\mathbf{k}'} e^{i\frac{\hbar}{c}\mathbf{k}'\mathbf{x}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \sum_{r'=1}^4 b_{*r',\mathbf{k}'}^\dagger \underline{e}_{or'}'^\dagger(\mathbf{k}') \cdot \\
&\quad \cdot \hbar \left( \begin{array}{c} \omega(\mathbf{k}) \sum_{r=1}^2 (\underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}}) - \\ - \omega(\mathbf{k}) \sum_{r=3}^4 (\underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}}) \end{array} \right) \\
&= \int_{(\Omega)} d\mathbf{x} \cdot \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \sum_{\mathbf{k}'} e^{i\frac{\hbar}{c}\mathbf{k}'\mathbf{x}} e^{-i\frac{\hbar}{c}\mathbf{k}\mathbf{x}} \cdot \\
&\quad \cdot \sum_{r'=1}^4 b_{*r',\mathbf{k}'}^\dagger \underline{e}_{or'}'^\dagger(\mathbf{k}') \left( \begin{array}{c} \sum_{r=1}^2 (\underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}}) - \\ - \sum_{r=3}^4 (\underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}}) \end{array} \right) \\
&= \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \sum_{\mathbf{k}'} \left( \int d\mathbf{x} \cdot e^{-i\frac{\hbar}{c}(\mathbf{k}-\mathbf{k}')\mathbf{x}} \right) \cdot \\
&\quad \cdot \sum_{r'=1}^4 b_{*r',\mathbf{k}'}^\dagger \left( \begin{array}{c} \sum_{r=1}^2 \left( \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}} \right) - \\ - \sum_{r=3}^4 \left( \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}} \right) \end{array} \right) .
\end{aligned}$$

Since

$$\int_{(\Omega)} d\mathbf{x} \cdot e^{-i\frac{\hbar}{c}(\mathbf{k}-\mathbf{k}')\mathbf{x}} = \left( \frac{2\pi c}{\hbar} \right)^3 \delta_{\mathbf{k},\mathbf{k}'}$$

then

$$\begin{aligned}
& \int_{(\Omega)} d\mathbf{x} \cdot \chi_*^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \\
&= \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \sum_{\mathbf{k}'} \left( \frac{2\pi c}{\hbar} \right)^3 \delta_{\mathbf{k},\mathbf{k}'} \cdot \\
&\quad \cdot \sum_{r'=1}^4 b_{*r',\mathbf{k}'}^\dagger \left( \begin{array}{c} \sum_{r=1}^2 \left( \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}} \right) - \\ - \sum_{r=3}^4 \left( \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{or}'(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{or'}'^\dagger(\mathbf{k}') \underline{e}_{*r}'(\mathbf{k}) b_{*r,\mathbf{k}} \right) \end{array} \right) .
\end{aligned}$$

In accordance with properties of  $\delta$ :

$$\begin{aligned}
& \int_{(\Omega)} d\mathbf{x} \cdot \chi_*^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \\
&= \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \left( \frac{2\pi c}{\hbar} \right)^3 .
\end{aligned}$$

$$\cdot \sum_{r'=1}^4 b_{*r',\mathbf{k}}^\dagger \left( \sum_{r=1}^2 \left( \underline{e}_{or'}^\dagger(\mathbf{k}) \underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{or'}^\dagger(\mathbf{k}) \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}} \right) - \sum_{r=3}^4 \left( \underline{e}_{or'}^\dagger(\mathbf{k}) \underline{e}'_{or}(\mathbf{k}) b_{or,\mathbf{k}} + \underline{e}_{or'}^\dagger(\mathbf{k}) \underline{e}'_{*r}(\mathbf{k}) b_{*r,\mathbf{k}} \right) \right).$$

Since

$$\begin{aligned} \underline{e}_{or'}^\dagger(\mathbf{k}') \underline{e}'_{or}(\mathbf{k}) &= \delta_{r,r'}, \\ \underline{e}_{or'}^\dagger(\mathbf{k}') \underline{e}'_{*r}(\mathbf{k}) &= 0 \end{aligned}$$

then

$$\begin{aligned} & \int_{(\Omega)} d\mathbf{x} \cdot \chi_*^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \\ &= \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \left( \frac{2\pi c}{h} \right)^3 \cdot \sum_{r'=1}^4 b_{*r',\mathbf{k}}^\dagger \left( \sum_{r=1}^2 (\delta_{r,r'} b_{or,\mathbf{k}} + 0 b_{*r,\mathbf{k}}) - \sum_{r=3}^4 (\delta_{r,r'} b_{or,\mathbf{k}} + 0 b_{*r,\mathbf{k}}) \right) \\ &= \left( \frac{2\pi c}{h} \right)^3 \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \cdot \sum_{r'=1}^4 b_{*r',\mathbf{k}}^\dagger \left( \sum_{r=1}^2 \delta_{r,r'} b_{or,\mathbf{k}} - \sum_{r=3}^4 \delta_{r,r'} b_{or,\mathbf{k}} \right) \\ &= \left( \frac{2\pi c}{h} \right)^3 \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \cdot \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \left( \sum_{r=1}^2 b_{*r,\mathbf{k}}^\dagger b_{or,\mathbf{k}} - \sum_{r=3}^4 b_{*r,\mathbf{k}}^\dagger b_{or,\mathbf{k}} \right). \end{aligned}$$

Therefore

$$\begin{aligned} & \int_{(\Omega)} d\mathbf{x} \cdot \chi_*^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \\ &= \left( \frac{2\pi c}{h} \right)^3 \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \left( \sum_{r=1}^2 b_{*r,\mathbf{k}}^\dagger b_{or,\mathbf{k}} - \sum_{r=3}^4 b_{*r,\mathbf{k}}^\dagger b_{or,\mathbf{k}} \right). \end{aligned}$$

Similarly you can calculate that

$$\begin{aligned} & \int d\mathbf{x} \cdot \chi_o^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \\ &= \left( \frac{2\pi c}{h} \right)^3 \sqrt{\frac{b - \sqrt{1 - a^2}}{b + \sqrt{1 - a^2}}} \sum_{\mathbf{k}} \hbar \omega(\mathbf{k}) \left( \sum_{r=1}^2 b_{or,\mathbf{k}}^\dagger b_{*r,\mathbf{k}} - \sum_{r=3}^4 b_{or,\mathbf{k}}^\dagger b_{*r,\mathbf{k}} \right). \end{aligned}$$

Since [5]

$$\tilde{\Psi}(t, \mathbf{p}) = \left( \frac{2\pi c}{h} \right)^3 \sum_{r=1}^4 c_r(t, \mathbf{p}) b_{r, \mathbf{p}}^\dagger \tilde{F}_0$$

and (10)

$$\left\{ b_{*r', \mathbf{k}'}^\dagger, b_{or, \mathbf{k}} \right\} = \hat{0}, \quad \left\{ b_{*r', \mathbf{k}'}^\dagger, b_{*r, \mathbf{k}}^\dagger \right\} = \hat{0}, \quad \left\{ b_{*r', \mathbf{k}'}^\dagger, b_{or, \mathbf{k}}^\dagger \right\} = \hat{0}.$$

then

$$b_{*r, \mathbf{k}}^\dagger b_{or, \mathbf{k}} \tilde{\Psi} = -b_{or, \mathbf{k}} \left( \frac{2\pi c}{h} \right)^3 \sum_{r=1}^4 c_r(t, \mathbf{p}) b_{*r, \mathbf{k}}^\dagger b_{r, \mathbf{p}}^\dagger \tilde{F}_0 = \tilde{0}.$$

Similarly

$$b_{or, \mathbf{k}}^\dagger b_{*r, \mathbf{k}} \tilde{\Psi} = \tilde{0}.$$

Hence

$$\boxed{\int_{(\Omega)} d\mathbf{x} \cdot \psi'^\dagger(\mathbf{x}) \hat{H}'_0 \psi'(\mathbf{x}) \Psi(t, \mathbf{x}_0) = \int_{(\Omega)} d\mathbf{x} \cdot \chi^\dagger(\mathbf{x}) \hat{H}'_0 \chi(\mathbf{x}) \Psi(t, \mathbf{x}_0).}$$

Thus, the function  $\psi'(\mathbf{x})$  can be substituted for the function  $\chi(\mathbf{x})$  in calculations of a probabilities evolution.

Let

$$\nu_{n_0, (s)}(\mathbf{k}) := \begin{bmatrix} (c + iq) e_{sL}(\mathbf{k}) \\ \vec{0}_2 \end{bmatrix}, \quad l_{n_0, (s)}(\mathbf{k}) := \begin{bmatrix} (a - ib) e_{sL}(\mathbf{k}) \\ e_{sR}(\mathbf{k}) \end{bmatrix}.$$

Hence from (6):

$$\cdot \left( \sqrt{\frac{2\pi n_0}{\sinh(2n_0\pi)}} \left( \begin{pmatrix} \cosh\left(\frac{h}{c} n_0 x_4\right) + \frac{e'_s(\mathbf{k}) = \frac{h}{2\pi c}}{\sinh\left(\frac{h}{c} n_0 x_4\right)} \\ + \sinh\left(\frac{h}{c} n_0 x_4\right) \end{pmatrix} (c + iq) e_{sL}(\mathbf{k}) + \right. \right. \\ \left. \left. + \begin{pmatrix} \cosh\left(\frac{h}{c} n_0 x_4\right) - \frac{e'_s(\mathbf{k}) = \frac{h}{2\pi c}}{\sinh\left(\frac{h}{c} n_0 x_4\right)} \\ - \sinh\left(\frac{h}{c} n_0 x_4\right) \end{pmatrix} \vec{0}_2 \right) + \exp\left(-i\frac{h}{c} (n_0 x_4)\right) l_{n_0, (s)}(\mathbf{k}) \right).$$

Therefore in the basis  $\mathbf{J}_{e\nu}$ :

$$\underline{e}'_s(\mathbf{k}) = \begin{bmatrix} \nu_{n_0, (s)}(\mathbf{k}) \\ l_{n_0, (s)}(\mathbf{k}) \end{bmatrix}.$$

Therefore from (12):

$$\chi(\mathbf{x}) = \sum_{\mathbf{k}} e^{-i\frac{h}{c} \mathbf{k} \cdot \mathbf{x}} \sum_{s=1}^4 \begin{bmatrix} \nu_{n_0, (s)}(\mathbf{k}) \\ l_{n_0, (s)}(\mathbf{k}) \end{bmatrix} b_{s, \mathbf{k}},$$



Let

$$\tilde{\nu}_{n_0}(\mathbf{x}) := \sum_{\mathbf{k}} e^{-i\frac{\mathbf{h}}{c}\mathbf{k}\mathbf{x}} \sum_{s=1}^2 \nu_{n_0,(s)}(\mathbf{k}) b_s(\mathbf{k}),$$

$$\tilde{l}_{n_0}(\mathbf{x}) := \sum_{\mathbf{k}} e^{-i\frac{\mathbf{h}}{c}\mathbf{k}\mathbf{x}} \sum_{s=1}^4 l_{n_0,(s)}(\mathbf{k}) b_s(\mathbf{k}).$$

Hence in the basis  $\mathbf{J}_{e\nu}$ :

$$\chi(\mathbf{x}) = \begin{bmatrix} \tilde{\nu}_{n_0}(\mathbf{x}) \\ \tilde{l}_{n_0}(\mathbf{x}) \end{bmatrix}. \quad (13)$$

Let:

$$\hat{H}_{l,0} := c \sum_{r=1}^3 \beta^{[r]} i \partial_r + \hbar n_0 (a \gamma^{[0]} - b \beta^{[4]}),$$

$$\hat{H}_{\nu,0} := c \sum_{r=1}^3 \beta^{[r]} i \partial_r + \hbar n_0 (a \gamma^{[0]} + b \beta^{[4]}),$$

$$\hat{H}_{\nu,l} := (c + iq) \begin{bmatrix} 0 & 0 & n_0 & 0 \\ 0 & 0 & 0 & n_0 \\ -n_0 & 0 & 0 & 0 \\ 0 & -n_0 & 0 & 0 \end{bmatrix},$$

$$\hat{H}_{l,\nu} := (c - iq) \begin{bmatrix} 0 & 0 & -n_0 & 0 \\ 0 & 0 & 0 & -n_0 \\ n_0 & 0 & 0 & 0 \\ 0 & n_0 & 0 & 0 \end{bmatrix}.$$

In that case in the basis  $\mathbf{J}_{e\nu}$ :

$$\hat{H}'_0 = \begin{bmatrix} \hat{H}_{\nu,0} & \hat{H}_{\nu,l} \\ \hat{H}_{l,\nu} & \hat{H}_{l,0} \end{bmatrix}.$$

Let

$$\hat{H}_{l,0}(\mathbf{k}) := \sum_{r=1}^3 \beta^{[r]} k_r + n_0 (a \gamma^{[0]} - b \beta^{[4]}),$$

$$\hat{H}_{\nu,0}(\mathbf{k}) := \sum_{r=1}^3 \beta^{[r]} k_r + n_0 (a \gamma^{[0]} + b \beta^{[4]}).$$

In that case

$$\hat{H}'_0(\mathbf{k}) = \begin{bmatrix} \hat{H}_{\nu,0}(\mathbf{k}) & \hat{H}_{\nu,l} \\ \hat{H}_{l,\nu} & \hat{H}_{l,0}(\mathbf{k}) \end{bmatrix}$$

An neutrino and it's lepton are tied by the follows equations:

$$\hat{H}_{\nu,0}(\mathbf{k}) \nu_{n_0,(s)}(\mathbf{k}) + \hat{H}_{\nu,l} l_{n_0,(s)}(\mathbf{k}) = \omega(\mathbf{k}) \nu_{n_0,(s)}(\mathbf{k})$$

for  $s \in \{1, 2\}$  and

$$\hat{H}_{\nu,0}(\mathbf{k}) \nu_{n_0,(s)}(\mathbf{k}) + \hat{H}_{\nu,l} l_{n_0,(s)}(\mathbf{k}) = -\omega(\mathbf{k}) \nu_{n_0,(s)}(\mathbf{k})$$

for  $s \in \{3, 4\}$ .

I suppose that such neutrino can fly 1.5 cm. [6] and give birth to it's leptons.

## References

- [1] G. Quznetsov, *Probabilistic Treatment of Gauge Theories*, in series Contemporary Fundamental Physics, ed. V. Dvoeglazov, Nova Sci. Publ., NY (2007), p. 98
- [2] Idem, p.p. 72, 75, 76
- [3] Idem, p.p. 98, 99
- [4] Idem, p. 77
- [5] Idem, p. 94
- [6] CDF Collaboration, Study of multi-muon events produced in pp collisions at  $\sqrt{s} = 1.96$  TeV, *preprint*  
<http://arxiv.org/abs/0810.5357>